



ONLINE ORGANIC GARDEN SUPPLIES
www.jamiesgardenshop.co.za

Jamie's Garden Shop

SOP for control of Powdery Mildew (PM) on Cannabis.

*If oil based pesticides have been used, make sure there is a 2 week waiting period before the first application of Bio-Sulphur.

Vegetative Period.

Spray once a week with Bio-Sulphur at 1-3ml per litre of water. If leaf damage occurs on sensitive cultivars reduce dosage amount, and increase the waiting period between sprays.

*Cut off any leaves with (PM) and dispose in the garbage.

Flower Period.

1 Week before flowering starts spray Bio-Sulphur at 1-3ml per litre of water.

At the very start of flower (1 week after your pre-flower spray) spray Bio-Sulphur at 1-3ml per litre of water.

1 Week after the start of flower spray your final spray of Bio-Sulphur at 1-3ml per litre of water.

1 Week after the final Bio-Sulphur Spray. Spray AQSF at 5-15ml per L water, combined with Bio-Tricho at 3-5ml per L water.

There after Spray AQSF at 5-15ml per L water, combined with Bio-Tricho at 3-5ml per L water. Once a week until week 7 of flower

*In high humidity areas spray Spray AQSF at 15ml per L water, combined with Bio-Tricho at 5ml per L water, twice a week until the last week of flower.